

JOURNAL OF ALGEBRA 45, 321-333 (1977)

## Induced Characters and Symplectic Sections

DAVID T. PRICE

*Department of Mathematics, Wheaton College, Wheaton, Illinois 60187**Communicated by Walter Feit*

Received February 17, 1976

## 1. INTRODUCTION

Let  $G$  be a finite group and  $\chi \in \text{Irr } G$  (the set of irreducible characters of  $G$ ). Let  $N$  be a normal subgroup of  $G$ . I. M. Isaacs noticed that when  $G/N$  is supersolvable, then there is a subgroup  $H$  of  $G$ , and  $\theta \in \text{Irr } H$  such that  $\theta^G = \chi$ , while  $\theta \mid N \in \text{Irr } N$ . This situation often can be exploited in inductive arguments (e.g., Step 2 in the proof of Theorem 3.2). In this paper we explore some aspects of such a configuration.

Let  $G$  be a solvable group with  $N \triangleleft G$ . When  $G/N$  is supersolvable, Dornhoff [1] found that restricting the Sylow subgroups of  $N$  to the modular or quaternion-free condition is the most general condition of the Sylow subgroups of  $N$  which guarantees that  $G$  is an  $M$ -group. When  $N$  satisfies the Dornhoff condition, we show in Theorem 4.1 the most general condition of  $G/N$  such that  $G$  is an  $M$ -group.

The results of Section 5 have been applied by Schacher and Seitz [9].

The notation in this paper is largely that of Huppert [2]. If  $H$  is a normal subgroup of the finite group  $G$ , and  $\theta \in \text{Irr } H$ , then the inertia group of  $\theta$  in  $G$  will be denoted  $I_G(\theta)$ .

Most of the results of this paper were contained in the author's doctoral dissertation, prepared under the supervision of G. Glauberman at the University of Chicago. During that period, the author was supported by a National Science Foundation Graduate Fellowship. The preparation of this paper was supported by a grant from the Wheaton College Alumni Association. Special thanks go to I. M. Isaacs for helpful conversations during this work.

## 2. PRELIMINARY RESULTS ABOUT CHARACTERS

In addition to the fundamental Frobenius Reciprocity Theorem and Clifford's theorem, the following lemmas will be useful.

LEMMA 2.1 ([4, Lemma 3.3]). *Let  $G$  be a finite group with subgroups  $A$*

and  $B$  such that  $G = AB$  and let  $\theta$  be any class function on  $A$ . Then  $\theta^G \mid B = (\theta \mid (A \cap B))^B$ .

The next lemma analyzes a situation which occurs frequently in this paper.

LEMMA 2.2. *Let  $N \triangleleft G$ , and  $\chi \in \text{Irr } G$ . Suppose that  $\chi \mid N \in \text{Irr } N$ . Suppose there is an  $H \leq G$  and  $\theta \in \text{Irr } H$  with  $\theta^G = \chi$ . Then:*

- (a)  $G = HN$ .
- (b)  $\theta \mid (H \cap N) \in \text{Irr}(H \cap N)$ .
- (c)  $\{\theta \mid (H \cap N)\}^N = \chi \mid N$ .

*Proof.* Now  $(\theta^{HN})^G = \theta^G = \chi$ , so  $\theta^{HN} \in \text{Irr } HN$ . By Frobenius reciprocity,  $\theta^{HN}$  is a constituent of  $\chi \mid HN$ . But  $\chi \mid N \in \text{Irr } N$ , so  $\chi \mid HN \in \text{Irr } HN$ . Thus,  $\theta^{HN} = \chi \mid HN$ . But then  $\theta^{HN}(1) = \chi(1) = \theta^G(1)$ . Therefore,  $HN = G$ . By Lemma 2.1,  $\chi \mid N = \theta^G \mid N = \{\theta \mid (H \cap N)\}^N$ . Then  $\{\theta \mid (H \cap N)\}^N \in \text{Irr } N$ , so  $\theta \mid (H \cap N)$  must be an irreducible character of  $H \cap N$ .

The next result is useful sharpening of Clifford's theorem [3, Proposition 3].

THEOREM 2.3. *Let  $G$  be a finite group,  $H|K$  be an elementary abelian chief section of  $G$ . Let  $\chi \in \text{Irr } H$  and suppose  $\chi$  is invariant in  $G$ . Then one of the following occurs.*

- (a)  $\chi \mid K \in \text{Irr } K$ .
- (b)  $\chi \mid K = e\theta$ , where  $\theta \in \text{Irr } K$  and  $e^2 = (H : K)$ .
- (c)  $\chi \mid K = \theta_1 + \cdots + \theta_{(H:K)}$ , and the  $\theta_i \in \text{Irr } K$  are distinct.

THEOREM 2.4. *Let  $G$  be a finite group,  $H|K$  an elementary abelian chief section of  $G$ . Let  $\theta \in \text{Irr } K$ , and suppose that  $G = \mathbf{I}_G(\theta)H$ . Then one of the following occurs.*

- (a)  $\theta^H = \chi_1 + \cdots + \chi_{(H:K)}$ , and the  $\chi_i \in \text{Irr } G$  are distinct.
- (b)  $\theta^H = e\chi$ , with  $\chi \in \text{Irr } H$  and  $e^2 = (H : K)$ .
- (c)  $\theta^H \in \text{Irr } H$ .

### 3. INDUCING CHARACTERS OVER A NORMAL SUBGROUP

This section presents the fundamental definition for the paper, and some consequences.

DEFINITION. Let  $\chi$  be an irreducible character of the finite group  $G$ , and  $N$  be a normal subgroup of  $G$ . Call  $\chi$  *induced over  $N$*  if there is a subgroup  $H$  of  $G$  containing  $N$  and irreducible character  $\theta$  of  $H$  so that  $\theta^G = \chi$  while  $\theta \mid N$  is irreducible. Such an  $H$  and  $\theta$  will be said to induce  $\chi$  over  $N$ .

*Remarks.* (1) I. M. Isaacs suggested the study of characters induced over a normal subgroup after he observed that if  $G/N$  is supersolvable, then every irreducible character of  $G$  is induced over  $N$ . Theorem 3.3 is a slight generalization of this observation.

(2) When  $G/N$  is abelian, R. L. Roth has also noted that every irreducible character of  $G$  is induced over  $N$  [8, Theorem 3.1].

(3) The irreducible character  $\chi$  of  $G$  is monomial if and only if  $\chi$  is induced over the identity subgroup of  $G$ . So the study of characters induced over a normal subgroup is a generalization of the study of monomial characters.

**LEMMA 3.1.** *Let  $\chi$  be an irreducible character of the finite group  $G$  which is induced over the normal subgroup  $N$  of  $G$ . Let  $K \triangleleft G$  be a subgroup of  $\text{Ker } \chi$ . Then  $\chi$  (as a character of  $G/K$ ) is induced over  $NK/K$ .*

The corollary to the next two theorems extends a result of Dornhoff [1].

**THEOREM 3.2.** *Let  $G$  be a finite group,  $N \triangleleft G$ . Suppose  $N$  is solvable, the odd Sylow subgroups of  $N$  are modular, and the Sylow 2-subgroups of  $N$  are quaternion-free. Let  $\chi \in \text{Irr } G$  and suppose that  $\chi$  is induced over  $N$ . Then  $\chi$  is a monomial character of  $G$ .*

*Proof.* By induction on  $|G|$ .

**Step 1.**  $\chi$  is faithful. Let  $K = \text{Ker } \chi$ . By Lemma 3.1,  $\chi$ , as a character of  $G/K$ , is induced over  $NK/K$ . Now  $NK/K \simeq N/(N \cap K)$  inherits the hypothesis on  $N$ , so, if  $K > 1$ , then by induction  $\chi$  is a monomial character of  $G/K$  and hence it is a monomial character of  $G$ .

**Step 2.**  $\chi|_N$  is irreducible. Let  $H \leq G$  with  $\theta \in \text{Irr } H$  induce  $\chi$  over  $N$ . Trivially,  $\theta$ , as a character of  $H$ , is induced by  $\theta$  over  $N$ . If  $H < G$ , then by induction,  $\theta$  is a monomial character of  $H$ , so  $\chi = \theta^G$  is a monomial character of  $G$ .

**Step 3.**  $\chi$  is primitive. Let  $H \leq G$ , and suppose, on the contrary, that there is  $\theta \in \text{Irr } H$  with  $\theta^G = \chi$ . Let  $L = H \cap N$ . Then  $L \triangleleft H$ , and since  $L \leq N$ , the assumptions on  $N$  are inherited by  $L$ . By Lemma 2.2(b),  $\theta|_L \in \text{Irr } L$ . Thus,  $\theta$  as a character of  $H$ , is induced over  $L$ . If  $H < G$ , then by induction  $\theta$  is a monomial character of  $H$ , so  $\chi = \theta^G$  is a monomial character of  $G$ .

**Step 4.** Let  $A \triangleleft G$ , with  $A$  abelian. Then  $A \leq \mathbf{Z}(G)$ , and  $A$  is cyclic. By Step 3 and Clifford's theorem,  $\chi|_A = \chi(1)\lambda$ , for some linear character  $\lambda$  of  $A$ . Since  $\chi$  is faithful,  $A \leq \mathbf{Z}(G)$ , and since  $A$  is represented by scalar matrices in a representation of  $G$  affording  $\chi$ ,  $A$  is cyclic.

**Step 5.**  $N$  is nilpotent. Let  $K$  be the smallest normal subgroup of  $N$  for

which  $N/K$  is nilpotent. If  $K > 1$ , let  $A$  be a nonidentity characteristic abelian subgroup of  $K$ . Then  $A \triangleleft G$ , so by Step 4,  $A \leq \mathbf{Z}(G)$ . But by Dornhoff [1],  $K \cap \mathbf{Z}(G) = 1$ , a contradiction.

Step 6.  $N$  is cyclic,  $\chi(1) = 1$ , and so  $\chi$  is monomial.

By Rigby [7], the odd Sylow subgroups of  $N$  are all cyclic, since they are modular, while the Sylow 2-subgroup of  $N$  is cyclic since it is quaternion-free. Hence  $N$  is the direct product of cyclic groups, and is itself cyclic. Now by Step 2,  $\chi \mid N \in \text{Irr } N$ , so  $\chi(1) = 1$ .

**THEOREM 3.3.** *Let  $G$  be a finite group, with  $N \triangleleft G$ . Suppose  $G/N$  is solvable, and for each  $H \geq N$ , the  $H$ -chief sections of  $H/N$  have odd rank. Let  $\chi \in \text{Irr } G$ . Then  $\chi$  is induced over  $N$ .*

*Proof.* By induction on  $(G : N)$ . Let  $\mathcal{S} = \{M \triangleleft G \mid N \leq M, \chi \mid M \notin \text{Irr } M\}$ . If  $\mathcal{S} = \emptyset$ , then  $\chi \mid N \in \text{Irr } N$ , and  $\chi$  is trivially induced over  $N$ . If  $\mathcal{S} \neq \emptyset$ , pick  $M \in \mathcal{S}$  with  $|M|$  as large as possible. Now  $M \neq G$  since  $\chi \in \text{Irr } G$ . Let  $K \triangleleft G$  so that  $K/M$  is a chief section of  $G$ . By the maximality of  $|M|$ ,  $\chi \mid K \in \text{Irr } K$ . Now  $|K/M| = p^d$  for some prime  $p$  and odd integer  $d$ . Inspect the three possibilities in Theorem 2.3. Since  $\chi \mid M \notin \text{Irr } M$ , (a) does not occur. Since  $d$  is odd, (b) does not occur. Hence

$$\chi \mid M = \sum_{i=1}^{p^d} \theta_i, \quad \theta_i \in \text{Irr } M.$$

Let  $H = \mathbf{I}_G(\theta_1)$ . By Clifford's theorem, there is  $\xi \in \text{Irr } H$ , such that  $\xi^G = \chi$ ,  $\xi \mid M = \theta_1$ , and  $(G : H) = p^d > 1$ . By induction  $\xi$  is induced over  $N$ , so  $\chi = \xi^G$  is induced over  $N$ .

**COROLLARY 3.4.** *Let  $G$  be a solvable group. Let  $N \triangleleft G$ . Suppose*

- (i) *for each  $N \leq H \leq G$ , all chief sections of  $H/N$  have odd rank;*
- (ii) *the odd Sylow subgroups of  $N$  are all modular, and the Sylow 2-subgroups of  $N$  are quaternion-free.*

*Then  $G$  is an  $M$ -group.*

**THEOREM 3.5.** *Let  $G$  be a finite group,  $N \triangleleft G$ . Let  $N \leq M \triangleleft G$  with  $(G : M)$  and  $(M : N)$  relatively prime. Let  $\chi \in \text{Irr } G$ , and let  $\theta$  be an irreducible constituent of  $\chi \mid M$ . Suppose  $\chi$  is induced over  $N$ . Then  $\chi$  is induced over  $M$  and  $\theta$  is induced over  $N$ .*

*Proof.* Let  $H \leq G$  with  $\xi \in \text{Irr } H$  induce  $\chi$  over  $N$ . Then  $\xi^{HM}$  is an irreducible constituent of  $\chi \mid HM$ . Hence  $\xi^{HM} \mid M$  is a sum of  $G$ -conjugates of  $\theta$ . Thus, replacing  $H$  and  $\xi$  by conjugates if necessary, we may assume that  $\theta$  is a con-

stituent of  $(\xi^{HM})|M$ . By Lemma 2.1,  $(\xi^{HM})|M = (\xi|(H \cap M))^M$ , and by Frobenius reciprocity,  $\xi|(H \cap M)$  is a constituent of  $\theta|(H \cap M)$ . Hence,  $\xi|N$  is an irreducible constituent of  $\theta|N$ .

Let  $\pi$  be the set of primes dividing  $(M:N)$ . Now,  $\chi(1)/\xi(1) = (G:H)$ . So

$$(\chi(1)/\xi(1))_\pi = (G:H)_\pi = (HM:H).$$

Also,  $\chi(1)/\theta(1)$  divides  $(G:M)$  [2, p. 570]. Thus  $\chi(1)/\theta(1)$  is a  $\pi'$ -number. Similarly,  $\theta(1)/\xi(1)$  is a  $\pi$ -number. But

$$\chi(1)/\xi(1) = \{\chi(1)/\theta(1)\}\{\theta(1)/\xi(1)\}.$$

Hence

$$\theta(1)/\xi(1) = \{\chi(1)/\xi(1)\}_\pi = (HM:H) = (M:H \cap M).$$

Now  $(\xi|(H \cap M))^M$  has  $\theta$  as a constituent, and

$$(\xi|(H \cap M))^M(1) = (M:H \cap M) \xi(1) = \theta(1).$$

Thus  $(\xi|(H \cap M))^M = \theta$ , and of course  $(\xi|(H \cap M))|N = \xi|N \in \text{Irr } N$ . Thus the character  $\xi|(H \cap M)$  of the subgroup  $H \cap M$  induces  $\theta$  over  $N$ . Similarly, since  $(\xi^{HM})|M = (\xi|(H \cap M))^M$  is an irreducible character of  $M$ , the character  $\xi^{HM}$  of the subgroup  $HM$  induces  $\chi$  over  $M$ .

Dade has given a method of embedding any solvable group in an  $M$ -group [2, Theorem V. 18.10]. The next lemma and theorem generalize his method to the situation studied in this section. Let  $G$  be a group, and  $H$  a permutation group on the symbol set  $\Omega$ . The wreath-product  $G \wr H$  of  $G$  with  $H$  is the set

$$\{(f, h) \mid h \in H, f \text{ mapping } \Omega \text{ into } G\},$$

together with the multiplication

$$(f_1, h_1)(f_2, h_2) = (g, h_1 h_2)$$

with  $g(i) = f_1(i)f_2(i^{h_1})$  for  $i \in \Omega$ .

**LEMMA 3.6.** *Let  $H$  be a finite group,  $N \triangleleft H$ , and suppose, for every  $\chi \in \text{Irr } H$ , that  $\chi$  is induced over  $N$ . Let  $Z$  be a cyclic group of prime order  $p$ , acting as a right regular permutation group on itself. Let  $G = H \wr Z$ . Let  $N_1 = \{(f, 1) \mid f: Z \rightarrow N\}$ . If  $\theta \in \text{Irr } G$ , then  $\theta$  is induced over  $N_1$ .*

The proof is similar to the one of Dade mentioned above.

**THEOREM 3.7.** *Let  $G$  be a finite group,  $N \triangleleft G$  with  $G/N$  solvable. Then there is a finite group  $G_1$  with normal subgroup  $N_1$  such that*

- (i)  $G \leq G_1$ .
- (ii)  $G \cap N_1 = N$ .
- (iii) Every  $\chi \in \text{Irr } G_1$  is induced over  $N_1$ .

#### 4. INDUCTIVE GROUPS

Let  $K$  be a finite group. Call  $K$  *inductive* if, whenever  $G$  is a finite group and  $\alpha$  is a homomorphism of  $G$  onto  $K$ , then every irreducible character of  $G$  is induced over  $\text{Ker } \alpha$ . In Theorem 3.3, we showed that if every chief section of every subgroup of the solvable group  $K$  has odd rank, then  $K$  is inductive. Theorem 4.1 gives two characterizations of inductive groups.

Let  $K$  be an inductive group. Every irreducible character of  $K$  is induced over the identity subgroup of  $K$ , and so  $K$  is an  $M$ -group. In particular,  $K$  is solvable.

**THEOREM 4.1.** *Let  $K$  be a finite solvable group. The following are equivalent.*

- (a)  $K$  is inductive.
- (b) Every subgroup of  $K$  has a covering group (for projective representations) which is an  $M$ -group.
- (c) No subgroup of  $K$  has a symplectic chief section. (A chief section of a solvable group is called symplectic if there is a nonsingular symplectic form on the section which is invariant under the inner automorphisms of the group.)

To illustrate the kind of  $M$ -group that is not inductive, let  $K = A_4$ , the alternating group on four letters. By Corollary 3.4, it is an  $M$ -group. Let  $G = SL(2, 3)$ . The faithful irreducible character  $\chi$  of  $G$  has degree 2, and cannot be induced over  $\mathbf{Z}(G)$ , since  $G/\mathbf{Z}(G) \simeq K$  has no subgroup of index 2. So  $K$  is not inductive. Also,  $G$  is a covering group of  $K$  and  $\chi$  is not monomial, so  $G$  is not an  $M$ -group, and  $K$  does not satisfy condition (b). Finally, the commutator map from  $\mathbf{O}_2(G)$  to  $\mathbf{Z}(G)$  induces a symplectic form on the minimal normal subgroup  $\mathbf{O}_2(K)$  of  $K$ , and  $K$  does not satisfy condition (c).

We now begin the proof of Theorem 4.1, and show that (c) implies something stronger than (b). Assume (c). Let  $H$  be a subgroup of  $K$ . Let  $F$  be any finite central extension by  $H$  (not just a covering group for  $H$ ), so there is  $C \leq \mathbf{Z}(F)$  with  $F/C \simeq H$ . If  $F$  is not an  $M$ -group, then some chief section  $A/B$  of some subgroup  $L$  of  $F$  is ramified [6, Theorem 1.2]. That means there is  $\theta \in \text{Irr } B$  which is invariant in  $L$  and  $\theta$  does not extend to an irreducible character of  $A$ . Thus,  $A/B$  admits a nonsingular  $L$ -invariant symplectic form [5, Theorem 2.7]. Define  $\alpha$  mapping  $L/C_L(A/B)$  onto  $LC/C_{LC}(AC/BC)$  by  $(gC_L(A/B))\alpha = gC_{LC}(AC/BC)$ . Since  $C \leq \mathbf{Z}(F)$  and  $A \not\leq \mathbf{Z}(F)$ , the map  $\alpha$  is an isomorphism. Now  $A/B$  and  $AC/BC$  are isomorphic  $LC$ -sections. Thus  $LC$  has the symplectic

section  $AC/BC$ , and then  $LC/C$  has the symplectic section  $\{AC/C\}/\{BC/C\}$ . But  $LC/C$  is isomorphic to a subgroup of  $K$ , contradicting (c).

We now prepare for the proof that (b) implies (a).

**PROPOSITION 4.2.** *Suppose that the finite solvable group  $H$  has a covering group  $H^*$  (for projective representations) which is an  $M$ -group. Let  $F$  be a finite extension by  $H$ , that is, there is a homomorphism  $\alpha$  of the finite group  $F$  onto  $H$ . Let  $N = \text{Ker } \alpha$ . Suppose  $\chi \in \text{Irr } F$  and  $\chi \upharpoonright N = e\theta$  for some integer  $e$  and  $\theta \in \text{Irr } N$ . Then  $\chi$  is induced over  $N$ .*

*Proof.* Since  $H^*$  is a covering group for  $H$ , there is a central subgroup  $C$  of  $H^*$ , and a mapping  $\rho$  of  $H$  into  $H^*$  so that  $h \rightarrow Ch^\rho$  is an isomorphism of  $H$  onto  $H^*/C$ . By Clifford's theorem [2, p. 567] and the properties of covering groups [2, p. 639], there is an ordinary matrix representation  $X$  of  $H^*$  of degree  $e$  and projective matrix representation  $Y$  of  $F$  of degree  $\theta(1)$  such that

$$T(f) = X(f^{\alpha\rho}) \otimes Y(f) \quad (f \in F),$$

is an ordinary matrix representation of  $F$  affording  $\chi$ . (The symbol  $\otimes$  denotes the Kronecker product of the matrices.)

By assumption  $H^*$  is an  $M$ -group. So there is a subgroup  $L^*$  of  $H^*$  of index  $e$  for which  $X \upharpoonright L^*$  has a linear constituent  $\lambda$ . Since  $C$  is represented by scalar matrices,  $C \leq L^*$ . Because  $X \upharpoonright L^*$  is completely reducible, we may conjugate the matrix representation  $X$  (if necessary), and assume that the matrices for  $X \upharpoonright L^*$  have  $\lambda$  in the  $(1, 1)$ -position, and all other entries in the first row and first column of  $X \upharpoonright L^*$  are zero. Let  $L = \{f \in F \mid f^{\alpha\rho} \in L^*\}$ . Let  $S(g) = \lambda(g^{\alpha\rho}) Y(g)$  for  $g \in L$ . From the shape of  $T(g)$ , we see that  $S$  is a constituent of  $T \upharpoonright L$  and  $S$  is an ordinary matrix representation of  $L$ . Let  $\phi$  be the character of  $L$  afforded by  $S$ . By construction  $\phi$  is a constituent of  $\chi \upharpoonright L$ . By Frobenius reciprocity,  $\chi$  is a constituent of  $\phi^F$ . But  $\phi^F(1) = (F:L)\phi(1) = (H^*:L^*)(\text{degree } Y) = e\theta(1) = \chi(1)$ . Hence  $\phi^F = \chi$ . Since  $\phi$  is a constituent of  $\chi \upharpoonright L$ , we have  $\phi \upharpoonright N$  is a constituent of  $\chi \upharpoonright N = e\theta$ . But  $\phi(1) = \theta(1)$ . Hence  $\phi \upharpoonright N = \theta$  and  $\phi$  induces  $\chi$  over  $N$ .

We now show that (b) implies (a) in Theorem 4.1. Assume (b). Let  $G$  be a finite group with normal subgroup  $N$  for which  $G/N \simeq K$ . Let  $\chi \in \text{Irr } G$ . Then  $\chi \upharpoonright N = e(\theta_1 + \cdots + \theta_t)$ , where  $e$  is a positive integer, and  $\theta_1, \dots, \theta_t$  are distinct irreducible characters of  $N$ . Let  $T = \text{I}_G(\theta_1)$ , the inertia group of  $\theta_1$ . Then there is a unique  $\xi \in \text{Irr } T$  for which  $\xi^G = \chi$  and  $\xi \upharpoonright N = e\theta_1$ . The group  $T$  with character  $\xi$  and normal subgroup  $N$  satisfy the hypothesis of Proposition 4.2. So there is a subgroup  $L$  of  $T$  and  $\phi \in \text{Irr } L$  for which  $\phi^T = \xi$  and  $\phi \upharpoonright N = \theta_1$ . Thus  $\phi^G = (\phi^T)^G = (\xi)^G = \chi$  and  $\phi$  induces  $\chi$  over  $N$ .

We finally show that an inductive group satisfies condition (c) of Theorem 4.1. Let  $K$  be an inductive group. Suppose, on the contrary, that there is a subgroup of  $K$  with a symplectic chief section. Take  $H \leq K$  of least possible order among

the subgroups with a symplectic chief section. Among the symplectic chief sections of  $H$ , choose one,  $M/L$ , such that  $M$  has the largest possible order. Then  $M/L$  is an elementary abelian  $p$ -group for some prime  $p$ . We fix these choices and notation for the rest of the argument. We shall eventually show that  $K$  is not inductive, a contradiction.

Let  $C = C_H(M/L)$ . Let  $D = O_p(H \text{ mod } C)$ . Since  $H/C$  has a nontrivial faithful irreducible module in characteristic  $p$ ,  $O_p(H \text{ mod } C) = C$ . Since  $H$  is solvable,  $D \neq C$ . Let  $S$  be a Hall  $p'$ -subgroup of  $D$ . Let  $A = LN_H(S)$ . By the Frattini argument,  $H = DN_H(S) = CA$ . Since  $M \triangleleft H$ , the product  $MA$  is a subgroup of  $H$ . Since  $H = CA$ , the elements of  $A$  cover the action of  $H$  on  $M/L$ . Hence  $M/L$  is a chief section of  $MA$ , and is symplectic. By the minimality of  $|H|$ , the subgroup  $MA$  cannot be proper, so  $H = MA$ .

Now consider  $M \cap A$ . Since  $M/L$  is abelian,  $M$  normalizes  $M \cap A$ . Of course  $A$  normalizes  $M \cap A$ . Hence  $M \cap A \triangleleft MA = H$ . Since  $M/L$  is a chief section of  $H$ ,  $M \cap A$  is either  $M$  or  $L$ . Suppose  $M \cap A = M$ , so that  $M \leq A$ . By Clifford's theorem,  $M = [M, D]L = [M, S]L$ . Hence,

$$M = [M, S]L \leq [A, S]L \leq SL.$$

But  $M/L$  is a  $p$ -group, while  $SL/L$  is a  $p'$ -group, a contradiction. Hence,  $M \cap A = L$ .

Let  $C_1 = C_A(M/L)$ . Note that  $C_1 \triangleleft MA = H$  and  $C_1 \cap M \leq A \cap M = L$ . Hence,  $C_1M/C_1$  is an  $H$ -chief section of  $H$  which is  $H$ -isomorphic to  $M/L$ . By the maximality of  $|M|$ , we have  $C_1M = M$  and  $C_1 = L$ . That is,  $C_A(M/L) = L$ .

Let  $H^* = H/L$ , and let  $M^*$  and  $A^*$  be the images of  $M$  and  $A$  in  $H^*$ . We summarize the structure of  $H^*$ . The subgroup  $M^*$  is a minimal normal subgroup of  $H^*$ . The subgroup  $A^*$  is a complement to  $M^*$  in  $H^*$ , and is faithfully represented by conjugation as a group of symplectic automorphisms of  $M^*$ .

If  $p$  is odd, let  $P$  be the extra-special group of exponent  $p$  and order  $p \mid M^* \mid$ . If  $p = 2$ , let  $P$  be the central product of an extra-special group of order  $2 \mid M^* \mid$  and a cyclic group of order 4. Let  $U$  be the group of automorphisms of  $P$  which are trivial on  $Z(P)$ . It is well known that  $I$ , the group of inner automorphisms of  $P$ , is a self-centralizing normal subgroup of  $U$ , and  $U/I$  induces (by conjugation) the full symplectic group on  $I$ . (Warning: When  $p = 2$ ,  $U$  need not be split over  $I$ .) Since  $M^*$  and  $I$  are isomorphic symplectic vector spaces, the linear group  $U/I$  has a subgroup  $B/I$  which is isomorphic to the group  $A^*$  of symplectic automorphisms of  $M^*$ . Let  $D = O_p(B \text{ mod } I)$ . Since  $B/I$  has a faithful irreducible module in characteristic  $p$ ,  $O_p(B \text{ mod } I) = B$ . Since  $B$  is solvable,  $D \neq I$ . Let  $S$  be a Hall  $p'$ -subgroup of  $D$ . Let  $N = N_B(S)$ . By the Frattini argument,  $B = IN$ . Since  $I$  is abelian,  $I \cap N \triangleleft I$ . Also,  $I \cap N \triangleleft N$  and hence  $I \cap N \triangleleft IN = B$ . Since  $I$  is an irreducible  $B/I$ -module,  $I \cap N$  is either 1 or  $I$ . If it is  $I$ , then  $I \leq N = N_B(S)$  and  $[I, S] \leq I \cap S = 1$ . This contradicts the fact that  $I$  is self-centralizing in  $U$ . Hence  $I \cap N = 1$ . Thus  $B$  is the semi-



direct product of  $I$  and  $N$ . By our choice of  $B$ , there is an isomorphism of  $H^*$  onto  $B$  which carries  $M^*$  to  $I$ .

So we may regard  $H^*$  as a group of automorphisms of  $P$  which are trivial on  $\mathbf{Z}(P)$  and in which the subgroup  $M^*$  induces the group of inner automorphisms on  $P$ .

Form the semidirect product  $PH^*$ . Then  $PM^*$  is extra-special if  $p$  is odd, and is the central product of an extra-special group with  $\mathbf{Z}(P)$  if  $p = 2$ . Let  $\theta$  be a faithful irreducible character of  $PM^*$ . Then  $\theta$  vanishes off of  $\mathbf{Z}(P)$  and thus  $\theta$  is invariant in  $PH^*$ . Let  $\lambda$  be the unique irreducible constituent of  $\theta \mid \mathbf{Z}(P)$ . Now  $\mathbf{Z}(P)H^* = \mathbf{Z}(P) \times H^*$  so consider the character  $\mu = \lambda \# 1_{H^*}$  of  $\mathbf{Z}(P)H^*$ . Let  $\chi$  be an irreducible constituent of  $\mu^{PH^*}$ . Then  $\mu$  is a constituent of  $\chi \mid \mathbf{Z}(P)H^*$  and hence  $\lambda$  is an irreducible constituent of  $\chi \mid \mathbf{Z}(P)$ . Since  $\theta$  is the only element of  $\text{Irr } PM^*$  which contains  $\lambda$  when restricted to  $\mathbf{Z}(P)$ , it is the only irreducible constituent of  $\chi \mid PM^*$ . Hence

$$\cdot \theta(1) \leq \chi(1) \leq \mu^{PH^*}(1) = (PH^*: \mathbf{Z}(P)H^*) = (P: \mathbf{Z}(P)) = \theta(1).$$

Thus  $\theta(1) = \chi(1)$  and  $\chi$  is an extension of  $\theta$  to  $PH^*$ .

We next argue that  $\chi$  is not induced over  $P$ . Now  $\chi \mid P = |M^*|^{1/2}\phi$  for some  $\phi \in \text{Irr } P$ . If  $\chi$  is induced over  $P$ , then there is  $J \leq PH^*$  with  $(PH^*: J) = |M^*|$ . By Lemma 2.2,  $PH^* = J(PM^*)$ . Hence,  $(PM^*: J \cap PM^*) = |M^*|^{1/2}$ . Since  $PM^*/P$  is abelian,  $J \cap PM^* \triangleleft PM^*$ . This contradicts the fact that  $PM^*/P$  is a chief section of  $PH^*$ .

Now form the semidirect product  $PH$  by using the natural homomorphism from  $H$  to  $H^*$  to define an action of  $H$  on  $P$ . Regard  $\chi$  as a character of  $PH$  having  $L$  in its kernel. Let  $T$  be a right transversal to  $H$  in  $G$  which contains 1. Let  $F = \{(\alpha, g) \mid \alpha: T \rightarrow P, g \in G\}$ . For  $(\alpha_1, g_1)$  and  $(\alpha_2, g_2)$  in  $F$ , define their product to be  $(\beta, g_1g_2)$ , where

$$\beta(t) = \alpha_1(t) \alpha_2(t')^{h'^{-1}}$$

for  $t \in T$  and  $tg_1 = h't'$  with  $h' \in H, t' \in T$ . Then  $F$  is the twisted wreath product  $P \wr_H G$  [2, pp. 99–100].

Let  $D = \{(\alpha, 1) \mid \alpha: T \rightarrow P, 1 \in G\}$ . Then  $D$  is isomorphic to a direct product of  $|T| = (G:H)$  copies of  $P$ .  $D$  is clearly normal in  $F$ . Also,  $F/D \simeq G$ . Let  $D_1 = \{(\alpha, 1) \mid (\alpha, 1) \in D, \alpha(1) = 1\}$ . Let  $S = \{(\alpha, h) \mid \alpha: T \rightarrow P, \alpha(t) = 1 \text{ if } t \neq 1, h \in H\}$ . Then  $S \simeq PH$ . Also,  $D_1 \cap S = 1$  and  $S$  normalizes  $D_1$ . Regard  $\chi$  as a character of  $D_1S$  having  $D_1$  in its kernel. Now  $\chi \mid D$  is homogeneous; let  $\xi$  be the irreducible constituent of  $\chi \mid D$ . Then  $D_1 \leq \text{Ker } \xi$  but  $D \cap S \not\leq \text{Ker } \xi$ . Now, if  $f \in F - D_1S$ , then  $f^{-1}(D \cap S)f \not\leq D_1$ . Hence,  $\xi^f \neq \xi$ . Thus, the inertia group of  $\xi$  lies in  $D_1S$ , so it must be  $D_1S$ . Let  $\psi = \chi^F$ . Let  $\beta$  be an irreducible constituent of  $\psi$ . By Frobenius reciprocity,  $\xi$  is a constituent of  $\beta \mid D$ . By Clifford's theorem, there is a unique  $\zeta \in \text{Irr}(\mathbf{I}_L(\xi))$  for which both  $\zeta \mid D$  has  $\xi$  as a constituent, and  $\beta$  is a constituent of  $\zeta^F$ . Furthermore,  $\zeta^F = \beta$ . But  $\chi$  satisfies

the conditions which characterize  $\zeta$ . Hence,  $\chi = \zeta$  and  $\chi^F = \beta$ . Thus,  $\chi^F = \psi$  is irreducible.

Now,  $G$  is an inductive group. So there is a subgroup  $J$  of  $F$  with  $D \leq J$ , and  $\alpha \in \text{Irr } J$  such that  $\alpha^F = \psi$  and  $\alpha|_D = \xi$ . Hence  $J \leq \mathbf{I}_L(\xi) = D_1 S$ . Consider  $\alpha^{D_1 S}$ . By the characterization of  $\zeta$  in the previous paragraph,  $\alpha^{D_1 S} = \zeta = \chi$ . Hence,  $\chi$  is induced over  $D$ . But  $D_1 \leq \text{Ker } \chi$ . So  $\chi$ , regarded as a character of  $S$ , is induced over  $D \cap S$ . But  $S \simeq PH$  and  $D \cap S \simeq P$ , contradicting the construction of  $\chi$  as a character of  $PH^*$  which was not induced over  $P$ . This contradiction shows that no subgroup of an inductive group has a symplectic chief section, and completes the proof of Theorem 4.1.

## 5. FORMATIONS OF INDUCTIVE GROUPS

We now consider the class  $\mathcal{F}$  of inductive groups. It is a formation (Theorem 5.1), but not a saturated formation (Proposition 5.2). The subformation  $\mathcal{O}$  of inductive groups of odd order is saturated, and we give a local definition, and an alternative description of  $\mathcal{O}$  (Theorem 5.3).

To show that a given group  $G$  is inductive, we employ Theorem 4.1, and show that no subgroup  $H$  of  $G$  has a symplectic chief section. We often argue that if  $H$  has a symplectic chief section, then some other group, which we are assuming to be inductive, also has a subgroup with a symplectic chief section. This contradicts Theorem 4.1.

**THEOREM 5.1.** *Let  $\mathcal{F}$  be the class of inductive groups.*

- (a) *If  $G \in \mathcal{F}$  and  $H$  is a subgroup of  $G$ , then  $H \in \mathcal{F}$ .*
- (b) *If  $G \in \mathcal{F}$  and  $H$  is a homomorphic image of  $G$ , then  $H \in \mathcal{F}$ .*
- (c) *If  $G_1$  and  $G_2$  are in  $\mathcal{F}$ , then  $G_1 \times G_2$  is in  $\mathcal{F}$ .*
- (d)  *$\mathcal{F}$  is a variety and a formation.*
- (e) *Let  $G$  be a finite group with  $G/\mathbf{Z}(G) \in \mathcal{F}$ . Then  $G \in \mathcal{F}$ .*

*Proof.* (a) By Theorem 4.1, it is sufficient to show that no subgroup  $K$  of  $H$  has a symplectic chief section. But if  $K$  has a symplectic chief section, then  $K$  is a subgroup of  $G$  with a symplectic chief section, contradicting Theorem 4.1 for  $G$ .

(b) Similar to (a).

(c) By Theorem 4.1, it is sufficient to show that no subgroup  $H$  of  $G_1 \times G_2$  has a symplectic chief section. Since  $G_1$  and  $G_2$  centralize each other in  $G_1 \times G_2$ , every chief section of  $H$  is  $H$ -isomorphic to an  $H$ -chief section of  $H/(H \cap G_1)$  or an  $H$ -chief section of  $H/(H \cap G_2)$ . But  $H/(H \cap G_1)$  (resp.  $H/(H \cap G_2)$ ) is isomorphic to a subgroup of  $G_2$  (resp.  $G_1$ ) and, by Theorem 4.1, has no symplectic chief section.

(d) This follows immediately from parts (a), (b), and (c) and the definition of a variety and a formation.

(e) It suffices to show that no subgroup  $H$  of  $G$  has a symplectic chief section. By the Jordan-Hölder theorem, every chief section of  $H$  is  $H$ -isomorphic to one obtained in a chief series which refines the normal series  $1 \leq H \cap \mathbf{Z}(G) \leq H$ . Since  $H \cap \mathbf{Z}(G) \leq \mathbf{Z}(H)$ , no chief section of  $H$  below  $H \cap \mathbf{Z}(G)$  is symplectic. A chief section of  $H$  above  $H \cap \mathbf{Z}(G)$  is (trivially) isomorphic to a section of  $H/(H \cap \mathbf{Z}(G))$  which is  $H$ -isomorphic to  $H\mathbf{Z}(G)/\mathbf{Z}(G)$ , a subgroup of  $G/\mathbf{Z}(G)$ . By Theorem 4.1,  $H\mathbf{Z}(G)/\mathbf{Z}(G)$  does not have a symplectic chief section.

PROPOSITION 5.2.  $\mathcal{F}$  is not a saturated formation.

*Proof.* Let  $p$  be a prime congruent to 1 mod 24, for example, take  $p = 73$ . Let  $F$  be the field of  $p$  elements. Let  $P = \{(u_1, u_2, u_3, u_4, u_5, u_6) \mid u_i \in F\}$  and define a multiplication of  $P$  by setting

$$\begin{aligned} & (u_1, u_2, u_3, u_4, u_5, u_6)(v_1, v_2, v_3, v_4, v_5, v_6) \\ &= (u_1 + v_1, u_2 + v_2, u_3 + v_3 + u_1v_2, u_4 + v_4, u_5 + v_5, u_6 + v_6 + u_4v_5). \end{aligned}$$

It is easy to verify that  $P$  is a group of order  $p^6$  and exponent  $p$ . The Frattini subgroup  $\Phi(P)$  is  $\{(0, 0, u_3, 0, 0, u_6) \mid u_3, u_6 \in F\}$ .

Let  $H = \langle x, t \mid x^3 = t^8 = 1, t^{-1}xt = x^{-1} \rangle$ , a group of order 24. Fix a primitive third root  $\omega$  and a primitive fourth root  $\rho$  of unity in  $F$ . Form the semidirect product  $G = PH$  by setting

$$x^{-1}(u_1, u_2, u_3, u_4, u_5, u_6)x = (\omega u_1, \omega u_2, \omega^2 u_3, \omega^2 u_4, \omega^2 u_5, \omega u_6)$$

and

$$t^{-1}(u_1, u_2, u_3, u_4, u_5, u_6)t = (u_4, u_5, u_6, \rho u_1, \rho u_2, -u_3),$$

for  $(u_1, u_2, u_3, u_4, u_5, u_6) \in P$ . It is easy to verify that these definitions extend to an action of  $H$  on  $P$ . Note that  $\Phi(P) \triangleleft G$ , then that  $\Phi(P) = \Phi(G)$ .

We now show that  $G/\Phi(G)$  is inductive. Since  $p \equiv 1 \pmod{24}$ , any noncyclic chief section of a subgroup of  $G/\Phi(G)$  must be in fact  $H$ -invariant, and be a subquotient  $V$  of  $P/\Phi(P)$ . Switch to additive notation on  $V$ . The element  $t$  induces a linear transformation on  $V$ , and the determinant of the action of  $t$  is  $-\rho$ . Since all symplectic transformations have determinant 1,  $V$  is not a symplectic chief section. By Theorem 4.1,  $G/\Phi(G)$  is inductive.

Now consider the minimal normal subgroup  $\Phi(G)$  of  $G$ . In additive notation, it is a two-dimensional vector space over  $F$ . The elements  $x$  and  $t$  induce linear transformations on  $\Phi(G)$  of determinant 1. Since the symplectic and special

linear groups coincide in dimension 2,  $\Phi(G)$  is a symplectic chief section of  $G$ , and by Theorem 4.1,  $G$  is not inductive. This group  $G$  shows that  $\mathcal{F}$  is not saturated.

The example constructed in the previous proof depended on special properties of the prime 2. We now consider the subformation  $\mathcal{O}$  of inductive groups of odd order. Schacher and Seitz [9] considered the formation  $\mathcal{J}$  of (solvable) groups of odd order in which every chief section of every subgroup has odd rank (compare Theorem 3.3). As is alluded to in the last paragraph of [9], the formations  $\mathcal{J}$  and  $\mathcal{O}$  are identical.

We define the local formations which will be used to induce  $\mathcal{O}$ .  $\mathcal{O}(2)$  is the empty formation. For  $p$  an odd prime,  $\mathcal{O}(p)$  is the formation of (solvable) groups  $H$  of odd order such that for every prime divisor  $q \in \pi(H) - \{p\}$ , the prime  $p$  has odd order in the multiplicative group of  $Z_q$ , the field of integers modulo  $q$ .

**THEOREM 5.3.** *The formations  $\mathcal{O}$  and  $\mathcal{J}$  are identical. The formation  $\mathcal{O}$  is locally induced by the  $\mathcal{O}(p)$ .*

*Proof.* Let  $\mathcal{L}$  be the formation locally induced by the  $\mathcal{O}(p)$ . We show  $\mathcal{J} \leq \mathcal{O} \leq \mathcal{L} \leq \mathcal{J}$ .

(1)  $\mathcal{J} \leq \mathcal{O}$ . This has been shown in Theorem 3.3.

(2)  $\mathcal{O} \leq \mathcal{L}$ . Let  $G \in \mathcal{O}$  and  $K/L$  be a chief section of  $G$ . Then  $K/L$  is an elementary abelian  $p$ -group for some odd prime  $p$ . Let  $C = C_G(K/L)$  and  $A = G/C$ . Let  $q \in \pi(A) - \{p\}$ . Take  $x \in G$  so that  $xC$  has order  $q$  in  $A$ . Since  $x$  does not centralize  $K/L$ , some  $\langle x \rangle$ -invariant irreducible subquotient  $K_0/L_0$  of  $K/L$  is not centralized by  $x$ . By Theorem 4.1 applied to the subgroup  $\langle x \rangle C$  of  $G$ ,  $K_0/L_0$  is not a symplectic  $\langle x \rangle$ -module. By [6, Lemma 2.2(a)],  $p$  has odd order in  $Z_q$ . Thus  $G \in \mathcal{L}$ .

(3)  $\mathcal{L} \leq \mathcal{J}$ . Let  $G \in \mathcal{L}$ . Since every subgroup of a group in  $\mathcal{L}$  is again in  $\mathcal{L}$ , we may assume, by induction on  $|G|$ , that every proper subgroup of  $G$  has all chief sections of odd rank. By [9, Lemma 4], every chief section of  $G$  has odd rank. Thus  $G \in \mathcal{J}$ .

## REFERENCES

1. L. DORNHOFF,  $M$ -groups and 2-groups, *Math. Z.* **100** (1967), 226–256.
2. B. HUPPERT, "Endliche Gruppen I," Springer, Berlin/Heidelberg/New York, 1967.
3. I. M. ISAACS, Fixed points and characters in groups with non-coprime operator groups, *Canad. J. Math.* **20** (1968), 1315–1320.
4. I. M. ISAACS, Extensions of group representations over nonalgebraically closed fields, *Trans. Amer. Math. Soc.* **141** (1969), 211–228.
5. I. M. ISAACS, Characters of solvable and symplectic groups, *Amer. J. Math.* **95** (1973), 594–635.
6. D. T. PRICE, Character ramification and  $M$ -groups, *Math. Z.* **130** (1973), 325–337.

7. J. F. RIGBY, Primitive linear groups containing a normal nilpotent subgroup larger than the centre of the group, *J. London Math. Soc.* **35** (1960), 389–400.
8. R. L. ROTH, A dual view of the Clifford theory of characters of finite groups, II, *Canad. J. Math.* **25** (1973), 1113–1119.
9. M. SCHACHER AND G. M. SEITZ,  $\pi$ -groups that are  $M$ -groups, *Math. Z.* **129** (1972), 43–48.